



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

YAP1-fusions in pediatric NF2-wildtype meningioma

Sievers, Philipp ; Chiang, Jason ; Schrimpf, Daniel ; Stichel, Damian ; Paramasivam, Nagarajan ; Sill, Martin ; et al ; Rushing, Elisabeth

DOI: <https://doi.org/10.1007/s00401-019-02095-9>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-178933>

Journal Article

Accepted Version

Originally published at:

Sievers, Philipp; Chiang, Jason; Schrimpf, Daniel; Stichel, Damian; Paramasivam, Nagarajan; Sill, Martin; et al; Rushing, Elisabeth (2019). YAP1-fusions in pediatric NF2-wildtype meningioma. *Acta Neuropathologica*, 139(1):215-218.

DOI: <https://doi.org/10.1007/s00401-019-02095-9>

***YAP1*-fusions in pediatric *NF2*-wildtype meningioma**

Philipp Sievers^{1,2}, Jason Chiang³, Daniel Schrimpf^{1,2}, Damian Stichel^{1,2}, Nagarajan Paramasivam⁴, Martin Sill^{5,6}, Tenzin Gayden^{7,8}, Belen Casalini^{1,2}, David E. Reuss^{1,2}, James Dalton³, Kristian W. Pajtler^{5,6,9}, Daniel Hänggi¹⁰, Christel Herold-Mende¹¹, Elisabeth Rushing¹², Andrey Korshunov^{1,2,5}, Christian Mawrin¹³, Michael Weller¹⁴, Matthias Schlesner⁴, Wolfgang Wick^{15,16}, Nada Jabado^{7,8}, David T.W. Jones^{5,17}, Stefan M. Pfister^{5,6,9}, Andreas von Deimling^{1,2}, David W. Ellison³, Felix Sahm^{1,2,5}

¹Department of Neuropathology, Institute of Pathology, University Hospital Heidelberg, Heidelberg, Germany

²Clinical Cooperation Unit Neuropathology, German Consortium for Translational Cancer Research (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany

³Department of Pathology, St. Jude Children's Research Hospital, Memphis, TN, USA

⁴Bioinformatics and Omics Data Analytics, German Cancer Research Center (DKFZ), Heidelberg, Germany

⁵Hopp Children's Cancer Center Heidelberg (KITZ), Heidelberg, Germany

⁶Division of Pediatric Neurooncology, German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany

⁷Department of Human Genetics, McGill University, Montreal, QC, H3A 1B1, Canada

⁸Department of Pediatrics, McGill University, and The Research Institute of the McGill University Health Center, Montreal, QC, H4A 3J1, Canada

⁹Department of Pediatric Oncology, Hematology, Immunology and Pulmonology, University Hospital Heidelberg, Heidelberg, Germany

¹⁰Department of Neurosurgery, University Medical Centre Mannheim, University of Heidelberg, Mannheim, Germany

¹¹Division of Experimental Neurosurgery, Department of Neurosurgery, University Hospital Heidelberg, Heidelberg, Germany

¹²Institute of Neuropathology, University Hospital and University of Zurich, Zurich, Switzerland

¹³Institute of Neuropathology, Otto-von-Guericke University, Magdeburg, Germany

¹⁴Department of Neurology, University Hospital and University of Zurich, Zurich, Switzerland

¹⁵Clinical Cooperation Unit Neurooncology, German Consortium for Translational Cancer Research (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany

¹⁶Department of Neurology and Neurooncology Program, National Center for Tumor Diseases, Heidelberg University Hospital, Heidelberg, Germany

¹⁷Pediatric Glioma Research Group, German Cancer Research Center (DKFZ), Heidelberg, Germany

Corresponding author:

Felix Sahm, MD, PhD

Department of Neuropathology

University Hospital Heidelberg

69120 Heidelberg, Germany

Fon: +49-6221 56-37886

Fax: +49-6221 56 4566

felix.sahm@med.uni-heidelberg.de

Key words: pediatric meningioma, molecular diagnostics, gene fusions, RNA sequencing, *YAP1-MAML2*, *YAP1-PYGO1*, *YAP1-LMO1*

Word count:

Main text 781, legend to figure in main manuscript 145

1 Meningioma is the most common primary central nervous system (CNS) tumor [8]. In
2 contrast to adulthood, meningiomas are rare among children and adolescents and frequently
3 (about 38%) occur in the context of tumor predisposition syndromes [12]. In line with the
4 frequent inactivation of *NF2* in adult meningiomas, neurofibromatosis type 2 is the most
5 common inherited syndrome predisposing to the early development of meningiomas, which
6 are often multiple. Other germline alterations predisposing to meningioma development are
7 *SMARCE1* [14] and *SUFU* mutations [1]. More recently identified drivers of meningiomas
8 include *AKT1/TRAF7*, *SMO*, *KLF4/TRAF7*, and *PIK3CA* mutations [3, 5].

9 The mutational underpinnings of sporadic pediatric meningioma have remained elusive to
10 date. We report in-frame gene rearrangements predicted to result in fusions involving *YAP1*
11 in nine meningiomas. We initially identified a *YAP1-MAML2* fusion by clinical RNA
12 sequencing in a four year old female patient with an intraventricular mass, histologically
13 compatible with meningioma. Subsequently, based on our database of >30,000 DNA
14 methylation profiles of brain tumors [4], including >1,000 meningiomas (among them about
15 102 pediatric meningiomas, defined as age of diagnosis equal or below 18 years) [13], and
16 corresponding copy number information, we additionally identified eight meningiomas with
17 structural alterations affecting chromosome 11q around the *YAP1* locus (Online Resources
18 Supplementary Fig. 2 and 3). All clustered with reference meningioma cases in t-SNE
19 analysis of DNA methylation data (Fig. 1a) and showed histological and
20 immunohistochemical features of meningioma (Online Resource Supplementary Fig. 1),
21 despite two being initially diagnosed as glioma. Interestingly, of the nine cases two were
22 classified as transitional and two as atypical meningiomas - both subtypes which often carry
23 *NF2* mutations. With the presented case number and incomplete information on few
24 samples, robust conclusions are, however, not derivable on subtype distribution. This
25 limitation also applies to localizations. The clinicopathological characteristics are summarized
26 in Table 1. RNA (seven samples) or exome (one sample) sequencing [15] revealed the
27 presence of *YAP1* fusions in all eight additional tumors (Fig. 1b-d and Online Resource
28 Supplementary Table 1).

29 Seven tumors harbored a rearrangement of *YAP1-MAML2* involving exons 1-5 (n=5) or only
30 in exon 1 (n=2) of *YAP1* (NM_001130145) and exon 2-5 of *MAML2* (NM_032427). *YAP1-*
31 *MAML2* fusions were verified by fluorescence *in situ* hybridization (FISH) performed in two
32 cases (Online Resource Supplementary Fig. 4). A *YAP1-PYGO1* fusion was seen in a single
33 case, containing exons 1-4 of *YAP1* and exons 2-3 of *PYGO1* (NM_015617). Additionally, a
34 *YAP1-LMO1* fusion was detected in another case involving exons 1-4 of *YAP1* and exons 2-
35 4 of *LMO1*. Of note, seven of the additional eight patients were children or adolescents,
36 whereas one patient was an adult.

37 *YAP1* is a transcriptional co-activator and downstream effector of the HIPPO pathway that
38 acts mainly through TEAD family transcription factors and regulates expression of genes
39 involved in cell proliferation and apoptosis [6, 7, 19, 20]. Deregulation of the HIPPO pathway
40 via overexpression of *YAP1*, leading to tumorigenesis is a frequent event in human
41 malignancies including meningiomas [2, 9]. Rearrangements involving the *YAP1* gene have
42 also recently been implicated as a driver in different types of cancer. Valouev et al. reported
43 an in-frame gene fusion between *YAP1* and *MAML2* in nasopharyngeal carcinomas [16]. A
44 similar fusion between *YAP1* and *MAMLD1* has been described in ependymoma [10, 11].
45 Both *MAML2* and *MAMLD1* are members of the Mastermind gene family and act as
46 transcriptional co-activators of NOTCH signaling [17]. The *YAP1-MAML2* rearrangement
47 combines the transcriptional activation domain of *MAML2* with the TEAD-binding domain of

YAP1, which likely results in NOTCH-independent co-activation of TEAD-mediated HIPPO signaling [17]. While *PYGO1* has been associated with different types of cancer, structural rearrangements including *YAP1* and *PYGO1* have not been reported to the best of our knowledge. Notably, *PYGO1* also functions as a transcriptional co-activator in the Wnt pathway. *LMO1* acts as a transcriptional regulator with a tumor-promoting activity, but its role in tumors has not been well studied.

These alterations seem to act as an alternative to *NF2* inactivation, since no *NF2* alterations were detected in the present cohort. The *NF2* gene product, the tumor suppressor merlin, functions upstream of the HIPPO pathway and there is growing evidence suggesting a functional link between *NF2*, *YAP1* and activation of the HIPPO pathway [2, 18]. In line, *YAP1* fusion positive meningiomas clustered closer to *NF2* mutant cases than other pediatric meningiomas (Online Resource Supplementary Fig. 5). This observation parallels the virtually mutually exclusive alterations of *NF2* and *YAP1* in ependymoma. Since DNA methylation correlates with cell-of-origin, the proximity of *YAP1*- and *NF2*-altered cases in the clustering might indicate more similarities of their precursor cells compared to *SMARCE1* mutant meningiomas. However, the lack of *AKT1*, *SMO*, or *KLF4/TRAFF7* mutant pediatric meningiomas and the few *SMARCE1* cases are limitations of this analysis.

Our findings identify *YAP1* fusions as a potential oncogenic driver in the development of meningiomas, predominantly in pediatric patients, and strengthen the hypothesis that deregulation of the HIPPO pathway is a central mechanism in meningioma tumorigenesis. Further studies in larger cohorts are needed to determine additional downstream functional consequences and a possible prognostic role of *YAP1* alterations in meningiomas.

Acknowledgements

We thank L. Dörner and H. Y. Nguyen for skillful technical assistance and the microarray unit of the DKFZ Genomics and Proteomics Core Facility for providing Illumina DNA methylation array-related services. This study was supported by the German Cancer Aid (70112956) and Else Kröner-Fresenius Stiftung (EKFS 2015_A60). FS is a fellow of the Else Kröner Excellence Program of the Else Kröner-Fresenius Stiftung (EKFS 2017_EKES.24).

Figure legends

Fig. 1 a) Unsupervised hierarchical clustering of DNA methylation profiles in nine *YAP1*-fused meningiomas (MNG_YP1) alongside 128 well-characterized CNS neoplasms and control tissue shown in a two-dimensional representation of pairwise sample correlations using the 15,000 most variant probes by t-distributed stochastic neighbor embedding (t-SNE) dimensionality reduction. Reference methylation classes: ependymoma, posterior fossa group A (EPN_PFA), ependymoma, posterior fossa group B (EPN_PFB), ependymoma, RELA fusion (EPN_RELA), ependymoma, YAP fusion (EPN_YAP), subependymoma, posterior fossa (EPN_PF_SE), ependymoma, spinal (EPN_SPINE), subependymoma, spinal (EPN_SPINE_SE), subependymoma, supratentorial (EPN_ST_SE), meningioma (MNG), chordoma (CHORDM), melanoma (MELN), schwannoma (SCHW) and control tissue white matter (CONTROL). Schematic of the *YAP1-MAML2* fusion involving exons 1-5 of *YAP1* and exons 2-5 of *MAML2* (**b**), the *YAP1-PYGO1* fusion involving exons 1-4 of *YAP1* and exons 2-3 of *MAML2* (**c**) and the *YAP1-LMO1* fusion involving exons 1-4 of *YAP1* and exons 2-4 of *LMO1* (**d**).

Table1 Clinicopathological characteristics of the *YAP1*-fused meningioma cohort

Case #	Age (years)	Sex	Tumor location	Initial Diagnosis	Genetic alteration
1	4	F	lateral ventricles, 3rd ventricle	MNG	<i>YAP1:MAML2</i>
2	3	M	temporal	PXA	<i>YAP1:PYGO1</i>
3	1	M	3rd ventricle, lateral ventricle	pHGG	<i>YAP1:MAML2</i>
4	2	M	skull base	MNG	<i>YAP1:MAML2</i>
5	36	M	optic nerve	MNG	<i>YAP1:MAML2</i>
6	8	F	skull base (supra-/infratentorial)	MNG	<i>YAP1:LMO1</i>
7	17	M	cavernous sinus	MNG	<i>YAP1:MAML2</i>
8	7	F	parietal	MNG	<i>YAP1:MAML2</i>
9	7	F	frontal	MNG	<i>YAP1:MAML2</i>

MNG – meningioma, PXA – pleomorphic xanthoastrocytoma, pHGG – pediatric high grade glioma, F – female, M – male, WHO grade is provided in Supplementary Table 2.

References

- 1 Aavikko M, Li SP, Saarinen S, Alhopuro P, Kaasinen E, Morgunova E, Li Y, Vesanen K, Smith MJ, Evans DGet al (2012) Loss of SUFU function in familial multiple meningioma. *Am J Hum Genet* 91: 520-526 Doi 10.1016/j.ajhg.2012.07.015
- 2 Baia GS, Caballero OL, Orr BA, Lal A, Ho JS, Cowdrey C, Tihan T, Mawrin C, Riggins GJ (2012) Yes-associated protein 1 is activated and functions as an oncogene in meningiomas. *Mol Cancer Res* 10: 904-913 Doi 10.1158/1541-7786.MCR-12-0116
- 3 Brastianos PK, Horowitz PM, Santagata S, Jones RT, McKenna A, Getz G, Ligon KL, Palescandolo E, Van Hummelen P, Ducar MDet al (2013) Genomic sequencing of meningiomas identifies oncogenic SMO and AKT1 mutations. *Nat Genet* 45: 285-289 Doi 10.1038/ng.2526
- 4 Capper D, Jones DTW, Sill M, Hovestadt V, Schrimpf D, Sturm D, Koelsche C, Sahm F, Chavez L, Reuss DEet al (2018) DNA methylation-based classification of central nervous system tumours. *Nature* 555: 469-474 Doi 10.1038/nature26000
- 5 Clark VE, Erson-Omay EZ, Serin A, Yin J, Cotney J, Ozduman K, Avsar T, Li J, Murray PB, Henegariu Oet al (2013) Genomic analysis of non-NF2 meningiomas reveals mutations in TRAF7, KLF4, AKT1, and SMO. *Science* 339: 1077-1080 Doi 10.1126/science.1233009
- 6 Levy D, Adamovich Y, Reuven N, Shaul Y (2008) Yap1 phosphorylation by c-Abl is a critical step in selective activation of proapoptotic genes in response to DNA damage. *Mol Cell* 29: 350-361 Doi 10.1016/j.molcel.2007.12.022
- 7 Liu-Chittenden Y, Huang B, Shim JS, Chen Q, Lee SJ, Anders RA, Liu JO, Pan D (2012) Genetic and pharmacological disruption of the TEAD-YAP complex suppresses the oncogenic activity of YAP. *Genes Dev* 26: 1300-1305 Doi 10.1101/gad.192856.112
- 8 Louis DN, Ohgaki H, Wiestler OD, Cavenee WK (2016) WHO Classification of Tumours of the Central Nervous System. IARC, City
- 9 Overholtzer M, Zhang J, Smolen GA, Muir B, Li W, Sgroi DC, Deng CX, Brugge JS, Haber DA (2006) Transforming properties of YAP, a candidate oncogene on the chromosome 11q22 amplicon. *Proc Natl Acad Sci U S A* 103: 12405-12410 Doi 10.1073/pnas.0605579103
- 10 Pajtler KW, Witt H, Sill M, Jones DT, Hovestadt V, Kratochwil F, Wani K, Tatevossian R, Punchihewa C, Johann Pet al (2015) Molecular Classification of Ependymal Tumors across All CNS Compartments, Histopathological Grades, and Age Groups. *Cancer Cell* 27: 728-743 Doi 10.1016/j.ccell.2015.04.002
- 11 Parker M, Mohankumar KM, Punchihewa C, Weinlich R, Dalton JD, Li Y, Lee R, Tatevossian RG, Phoenix TN, Thiruvengatam Ret al (2014) C11orf95-RELA fusions drive oncogenic NF-kappaB signalling in ependymoma. *Nature* 506: 451-455 Doi 10.1038/nature13109
- 12 Pathmanaban ON, Sadler KV, Kamaly-Asl ID, King AT, Rutherford SA, Hammerbeck-Ward C, McCabe MG, Kilday JP, Beetz C, Poplawski NKet al (2017) Association of Genetic Predisposition With Solitary Schwannoma or Meningioma in Children and Young Adults. *JAMA Neurol* 74: 1123-1129 Doi 10.1001/jamaneurol.2017.1406
- 13 Sahm F, Schrimpf D, Stichel D, Jones DTW, Hielscher T, Schefzyk S, Okonechnikov K, Koelsche C, Reuss DE, Capper Det al (2017) DNA methylation-based classification and grading system for meningioma: a multicentre, retrospective analysis. *Lancet Oncol* 18: 682-694 Doi 10.1016/S1470-2045(17)30155-9
- 14 Smith MJ, O'Sullivan J, Bhaskar SS, Hadfield KD, Poke G, Caird J, Sharif S, Eccles D, Fitzpatrick D, Rawluk Det al (2013) Loss-of-function mutations in SMARCE1 cause an inherited disorder of multiple spinal meningiomas. *Nat Genet* 45: 295-298 Doi 10.1038/ng.2552
- 15 Stichel D, Schrimpf D, Casalini B, Meyer J, Wefers AK, Sievers P, Korshunov A, Koelsche C, Reuss DE, Reinhardt Aet al (2019) Routine RNA sequencing of formalin-fixed paraffin-embedded specimens in neuropathology diagnostics identifies diagnostically and therapeutically relevant gene fusions. *Acta neuropathologica*: Doi 10.1007/s00401-019-02039-3
- 16 Valouev A, Weng Z, Sweeney RT, Varma S, Le QT, Kong C, Sidow A, West RB (2014) Discovery of recurrent structural variants in nasopharyngeal carcinoma. *Genome Res* 24: 300-309 Doi 10.1101/gr.156224.113
- 17 Wu L, Sun T, Kobayashi K, Gao P, Griffin JD (2002) Identification of a family of mastermind-like transcriptional coactivators for mammalian notch receptors. *Mol Cell Biol* 22: 7688-7700 Doi 10.1128/mcb.22.21.7688-7700.2002

- 18 Zhang N, Bai H, David KK, Dong J, Zheng Y, Cai J, Giovannini M, Liu P, Anders RA, Pan D (2010) The Merlin/NF2 tumor suppressor functions through the YAP oncoprotein to regulate tissue homeostasis in mammals. *Dev Cell* 19: 27-38 Doi 10.1016/j.devcel.2010.06.015
- 19 Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, Xie J, Ikenoue T, Yu J, Li Let al (2007) Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control. *Genes Dev* 21: 2747-2761 Doi 10.1101/gad.1602907
- 20 Zhao B, Ye X, Yu J, Li L, Li W, Li S, Yu J, Lin JD, Wang CY, Chinnaiyan AMet al (2008) TEAD mediates YAP-dependent gene induction and growth control. *Genes Dev* 22: 1962-1971 Doi 10.1101/gad.1664408



Suppl. material YAP1_new.pdf

Aktueller Pfad: [YAP1](#) / Suppl. material YAP1_new.pdf

Herunterladen(55.0 MB)

Supplementary materials and methods

Sample collection

Besides the index cases emerged in routine diagnostics, additional cases of *YAP1*-fused meningioma and reference cases were retrieved from the archives of the Departments of Neuropathology at the University Hospitals in Heidelberg, Magdeburg (both Germany) and Zürich (Switzerland), the Department of Pediatrics, McGill University, and The Research Institute of the McGill University Health Center, Montreal (Canada) as well as the Department of Pathology, St. Jude Children's Research Hospital, Memphis, TN (USA). Analysis of tissue and clinical data was performed in accordance with local ethical regulations.

Immunohistochemistry

Immunohistochemistry was performed on a Ventana BenchMark ULTRA Immunostainer (Ventana Medical Systems, Tucson, AZ, USA) for all cases with sufficient material (n=5). Antibodies were directed against: epithelial membrane antigen (EMA; Clone GP1.4, mouse monoclonal, dilution 1:1000, Thermo Fisher Scientific, Fremont, CA, USA), somatostatin receptor 2A (SSTR2A; SS-8000-RM, rabbit monoclonal, dilution 1:10, Biotrend, Cologne, Germany) and Ki-67 (clone MIB-1, mouse monoclonal, 1:100 dilution, Dako Agilent, Santa Clara, CA, USA).

Nucleic acid extraction, DNA methylation analysis and RNA/exome sequencing

Representative tumor tissue with highest tumor cell content was histologically identified and selected for nucleic acid extraction using local standard protocols. DNA methylation profiling of all samples was performed using the Infinium MethylationEPIC (850k) BeadChip (Illumina, San Diego, CA, USA) or Infinium HumanMethylation450 (450k) BeadChip (Illumina) array according to the manufacturer's instructions. Filtering and genome-wide copy number analyses were performed as previously described, using the 'conumee' package in R (<http://www.bioconductor.org>) [2]. RNA and DNA sequencing were performed on a NextSeq 500 (Illumina) as previously described [1]. Fusion discovery used defuse and arriba (<https://github.com/suhrig/arriba/>).

Fluorescence *in situ* hybridization (FISH)

FISH was performed using the YAP1 Break Apart Probe (Empire Genomics, Williamsville, NY, USA) on interphase nuclei on 5µm FFPE sections in two cases according to the standard procedures and manufacturer's instructions. The 5' centromeric and 3' telomeric probes are labeled with 5-carboxyl-x-rhodamine and 5-fluorescein, respectively.

